

The far infrared mobility edge of the QCD Dirac spectrum

Tamás G. Kovács

Eötvös Loránd University, Budapest, Hungary
and
Institute for Nuclear Research, Debrecen, Hungary




ELTE
EÖTVÖS LORÁND
UNIVERSITY



Bratislava, 08/05/2026

Spectral density of the lattice overlap Dirac operator for $T > T_c$




mobility edge at $\pm\lambda_c$

Peak at zero in the spectral density!

Edwards et al. PRD 61 (2000); Alexandru & Horvath, PRD 92 (2015); 2404.12298; Kaczmarek, Mazur, Sharma, PRD 104 (2021) 2021

Mobility edge vs. temperature

$N_f = 2 + 1$ lattice with physical staggered quarks




Giordano, TKG, Pittler, arXiv:2602.10921.

Are all the $|\lambda| < \lambda_c$ modes localized?

effective dimension across the spectrum

- Modes close to zero get delocalized!
- Is there another mobility edge?
- How to focus on $\lambda \rightarrow 0$?
- Need large volumes.



Meng, Sun, Alexandru, Horváth, Liu, Wang, Yang, JHEP 12 (2024) 101.

Instanton-based random matrix model of the spectral peak

- (Anti)instanton
→ zero eigenvalue of $D(A)$ with $(-)+$ chirality eigenmode
- High T :
large instantons “squeezed out” in the temporal direction
→ dilute gas of instantons and antiinstantons
- Zero modes exponentially localized:

$$\psi(r) \propto e^{-\pi Tr}$$


Instanton-antiinstanton pair

The Dirac operator in the subspace of zero modes


$$D(A) = \begin{pmatrix} 0 & iw \\ iw & 0 \end{pmatrix}$$

$$w \propto e^{-\pi Tr}$$

Spectrum of $D(A)$



Instanton and antiinstanton




Instanton-antiinstanton pair

The Dirac operator in the subspace of zero modes


$$D(A) = \begin{pmatrix} 0 & iw \\ iw & 0 \end{pmatrix}$$

$$w \propto e^{-\pi Tr}$$

Spectrum of $D(A)$



Instanton and antiinstanton




Instanton-antiinstanton pair

The Dirac operator in the subspace of zero modes


$$D(A) = \begin{pmatrix} 0 & iw \\ iw & 0 \end{pmatrix}$$

$$w \propto e^{-\pi Tr}$$

Spectrum of $D(A)$



Instanton and antiinstanton




Instanton-antiinstanton pair

The Dirac operator in the subspace of zero modes


$$D(A) = \begin{pmatrix} 0 & iw \\ iw & 0 \end{pmatrix}$$

$$w \propto e^{-\pi Tr}$$

Spectrum of $D(A)$




Instanton and antiinstanton




Spectrum of $D(A)$ in dilute gas of instantons

The Dirac operator in the subspace of zero modes

Spectrum of $D(A)$



Instantons and antiinstantons



n_i instantons n_a antiinstantons

→ $|n_i - n_a|$ exact zero modes + mixing near zero modes

Dirac operator in the subspace of zero modes (ZMZ)

Work by E.V. Shuryak, J.J.M. Verbaarschot, T. Schäfer (1990-2000)...

- Given n_i instantons, n_a antiinstantons in 3d box of size L^3
- Construct $(n_i + n_a) \times (n_i + n_a)$ matrix:

$$D = \begin{pmatrix} \overbrace{0}^{n_i} & \overbrace{iW}^{n_a} \\ iW^\dagger & 0 \end{pmatrix}$$

- $w_{ij} = A \cdot \exp(-\pi T \cdot r_{ij})$ r_{ij} is the distance of instanton i and antiinstanton j

Random matrix model of $D(A)$ in the zero mode zone

- How to choose instanton numbers (n_i, n_a) and locations?

- Quenched lattice $T > 1.05 T_c \rightarrow$ free instanton gas

Bonati et al. PRL 110 (2013); Vig R. & TGK, PRD 103 (2021)

- n_i and n_a independent identical Poisson-distributed

$$p(n_i, n_a) = e^{-\chi V} \cdot \frac{(\chi V/2)^{n_i}}{n_i!} \cdot \frac{(\chi V/2)^{n_a}}{n_a!}$$

χ is the topological susceptibility

- Locations random (uniform)


- $\rightarrow D(A)$ in quenched QCD $\Leftrightarrow$ ensemble of random matrices

Fit parameters to quenched lattice Dirac spectrum

$T = 1.1 T_c$ overlap Dirac spectrum

- Two parameters:
 - χ – topological susceptibility: from exact zero modes $\rightarrow \chi = \langle Q^2 \rangle / V$
 - A – prefactor of the exponential mixing between zero modes
- Fit A to distribution of Dirac eigenvalues (lowest eigenvalue)

$L = 2.4\text{fm}$ fit




Fit parameters to quenched lattice Dirac spectrum

$T = 1.1 T_c$ overlap Dirac spectrum

- Two parameters:
 - χ – topological susceptibility: from exact zero modes $\rightarrow \chi = \langle Q^2 \rangle / V$
 - A – prefactor of the exponential mixing between zero modes
- Fit A to distribution of Dirac eigenvalues (lowest eigenvalue)


$L = 2.4\text{fm}$ fit

$L = 3.5\text{fm}$ prediction



Spectral density



Lattice vs. random matrix model, $L = 3.5$ fm



Delocalization in the far infrared

$\chi_{top} \approx \chi_{top}(T_c)$ in quenched


$$I_{0.5} = \int_0^{0.5} \rho_{unf}(s) ds$$



Delocalization in the far infrared


$\chi_{top} \approx 4\chi_{top}(T_c)$ in quenched

$$I_{0.5} = \int_0^{0.5} \rho_{unf}(s) ds$$




Delocalization for different instanton densities

λ_c decreases with χ_{top}



The mobility edge versus χ_{top} in units of $\chi_{top}(T_c)$

- $\lambda_c \rightarrow 0$ as $\chi_{top} \rightarrow 0$
- χ_{top} controlled by T, m_q



Conclusions

- The QCD Dirac spectrum above T_c has a far infrared mobility edge
- $\lambda_{cIR} \rightarrow 0$ as $m_q \rightarrow 0$ and/or $T \rightarrow \infty$
- Consistent with exact constraints on the Dirac spectrum (see Matteo's talk)
- Spectral peak singular at zero, but power has nontrivial corrections
- The far infrared part of the spectrum drives $U(1)_A$ breaking above T_c .